


Gabazine

Cat. No.:	HY-103533
CAS No.:	104104-50-9
Molecular Formula:	C ₁₅ H ₁₈ BrN ₃ O ₃
Molecular Weight:	368.23
Target:	GABA Receptor
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling
Storage:	4°C, sealed storage, away from moisture
	* In solvent : -80°C, 2 years; -20°C, 1 year (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (271.57 mM; Need ultrasonic)
DMSO : ≥ 75 mg/mL (203.68 mM)
* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.7157 mL	13.5785 mL	27.1569 mL
	5 mM		0.5431 mL	2.7157 mL	5.4314 mL
	10 mM		0.2716 mL	1.3578 mL	2.7157 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: PBS
Solubility: 7.14 mg/mL (19.39 mM); Clear solution; Need ultrasonic and warming and heat to 60°C
2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (6.79 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (6.79 mM); Clear solution
4. Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (6.79 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Gabazine is a selective and competitive antagonist of GABA _A receptor, with an IC ₅₀ of ~0.2 μM for GABA receptor.
IC ₅₀ & Target	0.2 μM (GABA receptor) ^[1] .
In Vitro	Both bicuculline and Gabazine (SR 95531) have been characterized as competitive inhibitors of GABA binding to the GABA _A

receptor. Gabazine is more potent than bicuculline at blocking currents elicited by GABA, with an IC_{50} for currents elicited by 3 μ M GABA of $\sim 0.2 \mu$ M and a Hill coefficient of 1.0. Gabazine reduces the currents elicited by 10 μ M alphaxalone by $\sim 30\%$, for responses of receptors containing wildtype $\beta 2$ subunits. The concentration of Gabazine requires producing half the maximal block is $\sim 0.2 \mu$ M. Gabazine also could only produce a partial block of currents gated by 300 μ M pentobarbital. The maximal reduction, again, is $\sim 30\%$, and the concentration of Gabazine required to produce half the maximal block is $\sim 0.15 \mu$ M^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Nature. 2025 Sep 10.
- Nat Neurosci. 2023 May;26(5):751-764.
- Phytomedicine. 2022 Jan 29;98:153965.
- Elife. 2024 May 29.
- iScience. 2023 Mar 3;26(4):106322.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Ueno S, et al. Bicuculline and gabazine are allosteric inhibitors of channel opening of the GABAA receptor. J Neurosci. 1997 Jan 15;17(2):625-34.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite F, Monmouth Junction, NJ 08852, USA