

IPTG (≥ 98%)

Cat # NB-45-00111

Product Information

Code NB-45-00111

CAS number 367-93-1

Molecular Formula $C_9H_{18}O_5S$

Molecular Weight 238.3

Storage -20°C, stored under nitrogen

* In solvent : -80°C, 6 months; -20°C, 1 month (stored under nitrogen)

5g (NB-45-00111-5g); 5x5g (NB-45-00111-5x5g);

Sizes 25g (NB-45-00111-25g)

Note For research use only.



Solvent and solubility

In vitro:

H2O: 100 mg/mL (419.64 mM; Need ultrasonic)

DMSO : ≥ 60 mg/mL (251.78 mM)

* "≥" means soluble, but saturation unknown.

Preparing Stock solutions:

Solvent Concentration	1 mg	5 mg	10 mg
1 mM	4.1964 mL	20.9820 mL	41.9639 mL
5 mM	0.8393 mL	4.1964 mL	8.3928 mL
10 mM	0.4196 mL	2.0982 mL	4.1964 mL

Please refer to the solubility information to select the appropriate solvent.

In vivo:

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (8.73 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (8.73 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (8.73 mM); Clear solution

Biological activity

Description:

IPTG is a molecular mimic of allolactose, a lactose metabolite that triggers transcription of the lac operon, and it is therefore used to induce protein expression where the gene is under the control of the lac operator.



In Vitro:

IPTG uptake by E. coli can be independent of the action of lactose permease, since other transport pathways are also

involved. At low concentration, IPTG enters cells through lactose permease, but at high concentrations (typically used for

protein induction), IPTG can enter the cells independently of lactose permease.

For reference only

For Research Use Only. Not for Diagnostic or Therapeutic Use.