#### A0659

Leader in Biomolecular Solutions for Life Science

# ZO-1 Rabbit pAb

Catalog No.: A0659 91 Publications

**Basic Information** 

**Observed MW** 

**Calculated MW** 

Polyclonal Antibody

**Cross-Reactivity** 

Applications WB, IF/ICC, ELISA

195kDa

195kDa

Category

Human,Rat



#### Background

This gene encodes a member of the membrane-associated guanylate kinase (MAGUK) family of proteins, and acts as a tight junction adaptor protein that also regulates adherens junctions. Tight junctions regulate the movement of ions and macromolecules between endothelial and epithelial cells. The multidomain structure of this scaffold protein, including a postsynaptic density 95/disc-large/zona occludens (PDZ) domain, a Src homology (SH3) domain, a guanylate kinase (GuK) domain and unique (U) motifs all help to co-ordinate binding of transmembrane proteins, cytosolic proteins, and F-actin, which are required for tight junction function. Alternative splicing results in multiple transcript variants encoding different isoforms.

## **Recommended Dilutions**

WB	1:500 - 1:1000	Gene
		7082
IF/ICC	1:50 - 1:200	

е	ID			
-				

**Swiss Prot** 007157

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 18-273 of human TJP1 (NP\_003248.3).

Synonyms ZO-1

€

## **Product Information**

Source www.abclonal.com Rabbit


Isotype lgG

Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from Rat lung, using ZO-1 Rabbit pAb (A0659) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 45s.



Immunofluorescence analysis of MCF7 cells using ZO-1 Rabbit pAb (A0659) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.