

Mouse anti-CD30, clone Ber-H2 (Monoclonal)

Clone no. Ber-H2

MONOSAN

Product name	Mouse anti-CD30, clone Ber-H2 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:50-1:200)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1-k
Clonality	Monoclonal
Clone number	Ber-H2
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD30, clone Ber-H2 (Monoclonal)

Clone no. Ber-H2

MONOSAN

Additional info

The antibody detects a formalin-resistant epitope that is expressed by Reed-Sternberg cells in classic Hodgkin lymphoma, the majority of anaplastic large cell lymphomas, primary cutaneous CD30 positive T-cell lymphoproliferative disorders and in embryonal carcinomas. Occasionally diffuse large B-cell lymphoma stains with this antibody. This antibody also stains plasma cells in paraffin-embedded tissue as well as reactive immunoblasts. The staining pattern of anti-CD30 in lymphoma and embryonal carcinoma is different, with the former being membranous and exhibiting Golgi zone accentuation in location, and the latter being membranous only.

References

1. Schwarting R, et al., Blood 1989 (74):1678-1689
2. Fonatsch C, et al., Genomics 1992 (14):825-826
3. George DH et al. Am J Surg Pathol. 2003 Apr;27(4): 487-93
4. Hedvat CV et al. Hum Pathol. 2002 Oct;33(10): 968-74
5. Dabbs DJ. 4th Edition Saunders Elsevier. 2014; p702

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES