

Datasheet: NB-47-02254-100ML

Description:	DONKEY SERUM:Preservative Free
Name:	DONKEY SERUM
Format:	Preservative Free
Product Type:	Serum
Quantity:	100 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
Functional Assays	▪			
Immunoassay	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Form Serum - liquid

Preservative Stabilisers None present

Product Information **Donkey Serum** is a sterile filtered preparation of donkey serum prepared from donkey whole blood following clotting.

References

1. Mou, H. *et al.* (2013) Generation of Multipotent Foregut Stem Cells from Human Pluripotent Stem Cells [Cell Stem Cell. 10: 385-97.](#)
2. Barghorn, S. *et al.* (2005) Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease. [J Neurochem. 95: 834-47.](#)
3. Rashid, S.T. *et al.* (2010) Modeling inherited metabolic disorders of the liver using human induced pluripotent stem cells. [J Clin Invest. 120: 3127-36.](#)
4. McBride, P.A. *et al.* (1988) Immunostaining of scrapie cerebral amyloid plaques with antisera raised to scrapie-associated fibrils (SAF). [Neuropathol Appl Neurobiol. 14: 325-36.](#)
5. Vallier, L. *et al.* (2009) Early cell fate decisions of human embryonic stem cells and mouse epiblast stem cells are controlled by the same signalling pathways. [PLoS One. 4: e6082.](#)
6. Debertin, G. *et al.* (2015) Tyrosine hydroxylase positive perisomatic rings are formed around various amacrine cell types in the mammalian retina. [J Neurochem. 134 \(3\): 416-28.](#)
7. Pradillo, J.M. *et al.* (2017) Reparative effects of interleukin-1 receptor antagonist in young and aged/co-morbid rodents after cerebral ischemia. [Brain Behav Immun. 61: 117-26.](#)

Storage Store at -20°C only. Storage in frost free freezers is not recommended. Avoid repeated freezing

and thawing as this may denature the product. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch.
------------------	----------------------------------

Regulatory	For research purposes only
-------------------	----------------------------
